## The Myth & Reality of Treating Runners

Amol Saxena, DPM

Palo Alto Medical Foundation, Dept. of Sports Medicine, Palo Alto, CA, USA 650. 853-2943 | <u>HeySax@aol.com</u> | <u>www.amolsaxena.com</u> Podiatrist 1992, 2000 & 2004 Olympic Trials & Nike Oregon Project/Bay Area Track Club Fellowship Director, PAFMG

What is Sports Medicine?

Many Myths in Medicine: Sports Medicine no different!

Stretching: no scientific proof for prevention

10% rule

Rest, boot, eventually will get better

Let pain be your guide

Certain conditions like Plantar fasciitis and Achilles tendinopathy get better in a year

What I learned from treating Olympians

Sometimes surgery is better sooner

A lot of voodoo out there. REALITY:

- 1. Real rehab (myth of stretching)
- 2. ESWT/RSW (aka shock wave)
- 3. Alter-G treadmill
- 4. Surgery

30 pts in 2008 Olympic Track & Field Trials:

9 used orthoses

8 had surgery

5 had ESWT/RSW

20 used Alter-G Treadmill to continue or resume training

14 made the Olympic team

40 pts in 2012 Olympic T & F Trials

16 made Olympic teams

11 had "ESWT" (RSW)

7 had surgery

11 used orthoses

9 used Alter-G for their rehab

2016 Olympics: 15 Olympians

One Gold, One Silver, Two Bronze, one 4th, 5th, 6th & 7th, & four finalists

11 had surgery, 12 ESWT

43 in Olympic trials

Nike Oregon Project

"Salazar Phenomenon": get better quicker, don't dwell

The reality...

Body parts wear out: heart, eyes, prostate, uterus, etc.

So do tendons & joints!

MORAL: RATION YOUR RUNNING!

Some considerations for youth athletes

Yes, there are growing pains...growth plates are cartilage portion of bones that grow

These areas can get irritated ie: calacaneal apophysitis (aka "Sever's disease"), Osgood-Schlatter's

No permanent damage ever documented

OK to Run!

Some tips:

Running speed-work (intervals) in the AM is associated w more injuries

Concrete is harder than asphalt which is harder than dirt

The less number of steps/min is associated w over-striding and injury

Should have >80 steps (1 foot strike)/min – elite athletes have 90+ (count 1 foot for 30 secs & X 2)

"Terrible Toos/Twos"

Too much too soon

Two different changes ie inc of mileage and intensity

Avoid peer pressure (seen more in group training)

2018 HOKA OneOne SuperClinic- Dr. Amol Saxena

So what to tell runners?

Injury is part of any sport particularly when you do too much too soon

Run on softer surfaces

Increase gradually (use time)

Caution if injury history

And...

Cadence:Page 205 of Born to Run

Runner improved his speed by trying to match his cadence to a clip-on metronome that went 90 beats/min

The less number of steps/min is associated w over-striding and injury

Should have >80 steps (1 foot strike)/min – elite athletes have 90+ (count 1 foot for 30 secs & X 2)

Runner with Orthoses?

Myth: Flatfeet need motion control shoes

Another Myth: Hi-arched feet need a curved, "non-motion control shoe"

Reality: very few runners need motion controlled shoes

Only evidence for shoe recommendation is perceived comfort!

Reality: Radial Soundwave

"Radial Soundwave" (RSW):

3 Treatments 1 wk apart

**Plantar Fasciitis** 

Achilles Paratendinosis & insertional tendinoses,

"Shin splints" (medial tibial periostitis)

Patients can continue w sports

RCTs: the best evidence we have!

EARLY INTERVENTION?

Benefits of RSW

Continue to be active

Prospective study on 74 Achilles Tendinopathy patients shows > 75% improved after 1 yr (Saxena et al JFAS 2011)

2018 HOKA OneOne SuperClinic- Dr. Amol Saxena No anesthetic required, can still have surgery later Treatment of plantar fasciitis Stretching arch, night splint Arch supports or more stable shoes Rolling on frozen water bottle Next phase: injection, custom inserts, soundwave If all else fails, surgery (93% success rate) And by the way...Surgery for Plantar Fasciitis 93% successful in returning runners back to sport Saxena, Foot & Ankle Int 2006 Achilles tendinopathy Evaluate/change (higher heel)shoes? Avoid bent knee stretching Eccentric strengthening Ice bucket Inserts Soundwave All else fails: surgery 95+% successful Stress Fractures Relative rest: 1-12 wks from running May need casting, boot Ice, no anti-inflammatories Address biomechanics, weaknesses Check Vit D level! Bone stimulation & shockwave for some Surgery for certain types "POLICE" British J of Sports Med 2012 Protection

Optimal Loading (Alter-G)

lce

Compression (Support)

Elevation

My Pneumonic: "POLICED"

2017 Journal of Foot & Ankle Surgery

Protection (bracing, splint, tape etc)

Optimal Loading (antigravity TM etc)

Ice (can be with compression)

Core (strengthening)

Education (exercises, coaching, evidenced based)

Diet (Vitamin D, other factors)

Alter-G Treadmill: Faster RTA after Achilles Surg JFAS Saxena & Granot

What about prevention?

Stretching? Never been proven (can try roller, slower "dynamic" warm-up)

Strengthening? Correct muscles

Proper shoes: minimalist, maximalist

Proper form: "Pronation is not the evil, but over-striding is!"

Can see www.amolsaxena.com, www.smiweb.org, www.aapsm.org

"Everyone's an athlete, some just don't know it yet"- George Sheehan, MD

www.amolsaxena.com Thank You! 650 853 2943